```
Sibeck
Ukhorskiy
Takahashi
Fox
Wygant
Spence
Kletzing
Giles
Rowland
Hoffman
Doug described strategy
Paul goes through his presentation
            certain number of events / year with a fixed duration (1 day/event, 1 event /month) excel spreadsheet tracks regions
                        no tracking of two vehicles
set of parameters define critical region
                        another spreadsheet describes orbital elements for two-probe
                        trace orbit
                        B,L tracing for L and MLT where did L, MLT, Mlat ranges come from?
            question from Sasha:
L shell seems very low for EMIC, magnetosonic
                        Paul: Definitions need more work and refinement
            Wygant: need to add rows, e.g. substorm injections
Geoff: "Midnight sources" includes substorm injections, ring current injections, etc.
Wygant: don't decide based on how many rows are satisfied by an orbit
            Paul: this defines a trade space
Wygant: include some text that states that number of rows is not a criterion.
            wygam. Include some text that states that humber of rows is not a chehon.

Reeves: Each row should be treated independently, to show how changing orbit affects that particular row and likelihood of observing that type of event during the mission.

Obrien: Percentage represents time that either vehicle is in the given box.

Orbits defined co-orbital with 0.02 (120 km) apogee difference
            probability of observing at least two events by at least one spacecraft each, during the two year mission
Kletzing: Do you do ensembles of 2-year missions, to account for the random distribution of events
Obrien: calculates how long is it between visits to this box - how likely are we to be outside the box for a whole day
Necessary for programmatic reasons / funding reasons to do final analysis at APL
Linkage between Obrien and flowdown - takes into account 2-vehicle, revisit times
Xinlin: region definition focused on VLF waves. ULF waves not included.
             Sasha: ULF waves should be included as they may strongly drive the orbit (largest near magnetopause, driving apogee)
            Obrien: what about angle between line of nodes and line of apsides
Obrien: I have code that calls ONERA open-source library to do orbit tracing, B/L tracing, calculation of invariants (supports wide range of magnetic field models). Should I
```

Geoff: Look at ability of these to fill out the top couple of objectives.

Action items:

contribute these to APL?

Reeves define rows in OBrien table Sasha, Xinlin flesh out Approach 1.1, 1.2 Wygant approach 1.4 Rowland and Sibeck 1.3